

Exam 1
Environmental Science 218
Applied Hydrogeology

1. A large puddle that is 0.05m thick and 20 m on each side goes from being a liquid at 20 degrees C to being ice at -5 degrees C. How much heat must be removed from the water for this to happen.

We calculate the mass and then the heat removed to go from water at 20 to water at 0, water at 0 to ice at 0 and ice at 0 to ice at -5.

$$m = \rho V = 1 \frac{g}{cm^3} \cdot (5cm \cdot 2000cm \cdot 2000cm) = 2 \times 10^7 g$$

$$Q = mc\Delta T = 2 \times 10^7 g \cdot \frac{1cal}{g^\circ C} \cdot (0 - 20) = -4 \times 10^8 cal$$

$$Q = -m L = -2 \times 10^7 g \cdot 79.7 \frac{cal}{g} = -1.594 \times 10^9 cal$$

$$Q = mc\Delta T = 2 \times 10^7 g \cdot \frac{0.5cal}{g^\circ C} \cdot (-5 - 0) = -5 \times 10^7 cal$$


$$Q = -4 \times 10^8 cal - 1.594 \times 10^9 cal - 5 \times 10^7 cal = -2.044 \times 10^9 cal$$

2. A stream hydrograph is shown below.

a) Find t_1 from the graph and use it to compute the total potential groundwater recharge via the baseflow method.


b) If the recession were to be interrupted at time 100 days, what is the remaining potential baseflow?

Stream Hydrograph



3. Consider the stream hydrograph below. Use the Rorabaugh method to find the total recharge. Recall that the steps are:

- Find t_1 for the first recession.
- Use t_1 to find the critical time t_c .
- Locate the time t_c past the peak for B.
- Extrapolate using the lines given to find Q_A and Q_B for the time t_c past the peak
Find the recharge



4. Consider the basin map shown at the end of the test.
 - a) Compute the effective uniform depth of precipitation using the simple average
 - b) Draw a Thiessen diagram and explain how to use the areas to find the EUD (**label areas but do not measure them**).
 - c) Draw an Isohyetal map and explain how to use the areas to find the EUD (**label the areas but do not measure them**).

5. A V-notch weir is used to measure the flow of a stream. Explain how this is done. If the height of the water is 0.25 ft, what is the flow?

6. The hydrologic equation is


$$\Delta S = I - O$$

$\Delta S = \text{Change in storage}$

$I = \text{Inflow}$

$O = \text{Outflow}$

Consider the Basin below:



Inflow	Outflow	Change in Storage
Stream A: 10 cu m/s	Evaporation=30cm/yr	0
Stream B: 30 cu m/s	Groundwater=?	
Stream C: 20 cu m/s		
Stream D: 5 cu m/s		
Direct Precip: EUD =20cm/yr		
Groundwater=?		

- What is the net inflow due to streams for a year?
- What is the net inflow due to direct precipitation for a year?
- What is the net outflow due to Evaporation for a year?
- If the change in storage is zero, what must the net groundwater flow be?

You may find the following helpful: #sec/yr = 3.15×10^7

Conversions, equations, etc.

$$\rho_{\text{water}} = 1 \text{ g/cm}^3$$

$Q = 2.54 h^{3/2}$ for a triangular weir. h measured in ft, Q in cu ft/s

$$c_{\text{water}} = 1 \frac{\text{cal}}{\text{g} \cdot ^\circ\text{C}}$$

$$c_{\text{ice}} = 0.5 \frac{\text{cal}}{\text{g} \cdot ^\circ\text{C}}$$

$$L_{\text{fusion}} = 79.7 \frac{\text{cal}}{\text{g}}$$

$$V_{ip} = \frac{Q_0 t_1}{2.3026} \quad (\text{potential groundwater dischg})$$

$$V_t = \frac{V_{ip}}{10^{(t/t_1)}} \quad (\text{volume of dischg remaining at time } t)$$

$$G = \frac{2(Q_B - Q_A)t_1}{2.3026} \quad (\text{Rorabaugh Method})$$

$$t_c = .2144 t_1$$

